

PHYS 813: Statistical Mechanics, Assignment 7

Due 4/24/08

1. A surface with N_0 adsorption centers has $N < N_0$ gas molecules adsorbed on it, at most one molecule per center. Assume that the canonical partition function for a single adsorbed molecule is known and denote it by $a(T)$. Neglect interactions between adsorbed molecules. Find the chemical potential of the system:
 - (a) using the canonical partition function;
 - (b) using the grand canonical partition function. In this case N can vary from 0 to N_0 .
2. As a continuation of the previous problem, consider the (ideal) gas which is in equilibrium with the surface. Find the pressure of the gas as a function of the fraction of average number of occupied sites and of temperature.
3. For a single electron in a magnetic field \mathbf{B} , the Hamiltonian is

$$\hat{H} = -\mu_B \hat{\boldsymbol{\sigma}} \cdot \mathbf{B}$$

where μ_B is a constant and $\hat{\boldsymbol{\sigma}}$ is the spin operator. Choose the z axis to be directed along \mathbf{B} and find the expression for the canonical density matrix elements ρ_{mn} in the representation in which

- (a) $\hat{\sigma}_z$ is diagonal;
- (b) $\hat{\sigma}_x$ is diagonal.

Next, calculate the average value of $\hat{\sigma}_z$ in both representations.

4. A system with two energy levels is populated by N distinguishable noninteracting particles at temperature T with occupations determined by the canonical distribution.
 - (a) Find the average energy per particle.
 - (b) Find the behaviour of this energy as $T \rightarrow 0$ and $T \rightarrow \infty$.
 - (c) Find the specific heat to the system.
 - (d) Find the behaviour of the specific heat as $T \rightarrow 0$ and $T \rightarrow \infty$. Interpret the obtained results.
5. Consider a dilute, noninteracting gas of N distinguishable diatomic molecules of mass m . Each molecule is a rigid rotor with $(2J + 1)$ -degenerate energy levels $\epsilon_J = \hbar^2 J(J + 1)/2I$, where I is the moment of inertia. Start from the general expression for the canonical partition function $Q_N(V, T) = \sum_n e^{-E_n/kT}$ where E_n is the total energy of N -molecule system. Calculate then the average energy, the specific heat at constant volume, and the entropy of the system in the limit $kT \gg \hbar^2/2I$. You may use without proof the partition function for the ideal monoatomic gas $Q_N^{\text{ideal}}(V, T) = [V(2\pi mkT)^{3/2}/h^3]^N$. *Hint:* In the assumed limit, $\sum_n f(n) = \int_0^\infty f(x)dx$.

6. Use the Debye model to calculate the internal energy and heat capacity of a one-dimensional atomic solid with length L for both high and low temperature. Assume periodic boundary conditions. The Debye model treats the solid as a set of coupled harmonic oscillators and approximates the unknown normal modes of the system as plane waves propagating with the velocity of sound.
7. Consider a free particle in a box with periodic boundary conditions, in the momentum representation. The particle is in equilibrium with a heat bath and therefore is described by the canonical distribution. Denote the eigenfunctions of the Hamiltonian as

$$\phi_{\mathbf{k}}(\mathbf{r}) = \frac{1}{L^{3/2}} e^{i\mathbf{k}\cdot\mathbf{r}}$$

- (a) Evaluate matrix elements of $e^{-\beta\hat{H}}$ in the basis $\phi_{\mathbf{k}}(\mathbf{r})$.
- (b) Find the canonical partition function in terms of L and $\lambda = h/\sqrt{2\pi mkT}$.
- (c) Find the density operator $\hat{\rho}$ in the basis $\phi_{\mathbf{k}}(\mathbf{r})$.
- (d) Calculate the average value of \hat{H} as $\text{Tr}(\hat{\rho}\hat{H})$. Express your answer in terms of L , λ , and T .